

Intracranial chondrosarcoma: review of the literature and report of 15 cases

Arthur G G C Korten, Hans J W ter Berg, Geert H Spincemaille, Ronald T van der Laan, Antoinet M Van de Wel

Abstract

The available data in the literature (177 cases), two current clinical patients, and cases which occurred in The Netherlands (13) were reviewed concerning the clinical presentation, pathological features, radiological data, and treatment options of chondrosarcoma of the cranial base.

The mean age of patients was 37 years, the male/female ratio 1: 1.1. The most frequent complaints were diplopia with oculomotor disorders (51%), headache (31%), and decreased hearing, dizziness, and tinnitus with statoacoustic dysfunction (21%). The mean duration of symptoms before diagnosis was 27 months.

The chondrosarcomas were located in the petrosal bone in 37% (47 cases), in the occipital bone and clivus in 23% (30 cases), in the sphenoid bone in 20% (25 cases) and to a lesser extent in frontal, ethmoidal, and parietal bones (14%). In 6% (eight cases) the primary location was in dural tissue. Radiological examinations showed bone destruction and variable calcification (CT), involvement of neuronal and vascular structures (MRI), and mostly hypovascularity on angiography. On histological examination 51% of tumours were classified as grade I, 11% grade II, 30% mesenchymal, and 8% myxoid. The mesenchymal type was the most malignant as illustrated by a strong tendency to intradural and cerebral growth and possibly occurrence in younger age groups. The treatment of choice until recently was surgery because of the critical location and local aggressive nature. Regrowth of tumour after surgery occurred in 53% of the patients (average after 32 months). Charged particle irradiation gave a five year survival of 83-94% and a local control rate of 78%-91%. Both in surgery and radiotherapy there is treatment related morbidity and mortality that should be considered when offering these therapies.

Recent promising results imply that charged particle radiotherapy, in combination with surgery, may be the therapeutic choice of the future.

(J Neurol Neurosurg Psychiatry 1998;65:88-92)

Keywords: chondrosarcoma; cranial base neoplasma

represents 0.15% of all cranial space occupying lesions and 6% of all skull base tumours.¹ It is assumed that chondrosarcomas originate from remnants of embryonal cartilage or from metaplasia of meningeal fibroblasts.^{2,3}

To the best of our knowledge, the last detailed review of the literature disclosed only 50 cases until 1985.⁴ This led us to review the available literature since 1985 in detail. This report also concerns two of our own clinical patients and 13 other patients treated in different hospitals throughout The Netherlands. The clinical history, pathological features, radiological findings, and treatment options are discussed. From these data we give an overview of the current diagnostic procedures and therapeutic options of intracranial chondrosarcoma.

Methods

In addition to the study by Hassounah *et al*,⁴ analysis of the literature from 1985 disclosed 127 cases.⁴⁻³⁷ Our own two clinical patients and 13 other cases retrieved from the files of the Dutch Committee on Bone Tumours, University Hospital Leiden, and the reported cases since 1985 along with the 50 cases from the review of Hassounah *et al*⁴ were analysed. Table 1 shows summarised data of the 15 Dutch patients.

Patients with chondrosarcoma from the nasopharynx or the paranasal sinus and extending into the skull base and chondrosarcoma as part of the syndromal diseases Maffucci or Ollier were excluded. The 27 patients of the series of Castro *et al* were also excluded because of paucity of detailed information.³⁸

Results

The sex ratio was 1:1.1. In 16 cases, the sex of the patient was not reported. The average age was 37 years (range 3 months-76 years). Age was not reported in 67 cases.

Signs and symptoms at initial presentation were described in 67 patients (table 2). The median time period between initial symptoms and moment of diagnosis was 15 months (ranging from 1 month to 144 months). This information was available in only 36 cases.

Over the years, different radiological evaluation methods have been used. Plain skull radiography, CT, MRI, and angiography were the most often used diagnostic tools. On T1 weighted MRI, chondrosarcomas had a low to intermediate signal intensity and were isointense or hypointense to grey matter. On proton density and T2 weighted images, they had high signal intensity and were hyperintense to grey matter.³⁹

Department of
Neurology and
Department of
Radiology,
Maaslandziekenhuis,
Sittard,
The Netherlands
A G G C Korten
H J W ter Berg
R T van der Laan

Department of
Neurosurgery,
Academic Hospital,
Maastricht,
The Netherlands
G H Spincemaille

Radiotherapy Institute
"Limburg", Heerlen,
The Netherlands
A M Van de Wel

Correspondence to:
Dr Arthur GGC Korten,
Maaslandziekenhuis, PO Box
5500, 6130 MB Sittard,
The Netherlands.

Received 7 January 1997 and
in final revised form
5 January 1998
Accepted 14 January 1998

Tumours originating from bone at the base of skull are rare. One such tumour, the chondrosarcoma, the most malignant cartilage tumour,

Table 1 Characteristics of 15 Dutch patients

No	Age/sex	Location	Pathology	Size	Symptoms/signs
1	31/M	PS; extension in pons and in Scav; destruction of SSph	Grade I	2½×3 cm	Headache 2 y; 1 y diplopia; sensory disturbance OD. Ptosis / VI dysfunction OD; cornea reflex R<L; sensory disturbance V branch 1 and 2 R
2	26/F	FM; SS	Grade I		Pain face 1 y; paraesthesias. V dysfunction L; possibly VII dysfunction L
3	18/F	PS; extension in clivus and SCav; petroclival junction	Grade II	1×2 cm	Several months diplopia on looking L; headache. VI dysfunction L
4	15/F	PS; extension in SSph	Grade I	3×2×2 cm	1 y Diplopia; headache. Dysfunction III L
5	23/M	PS; Clivus; SCav	Grade II		1 y Intermittent frontoparietal headache; diplopia horizontally. VI dysfunction R
6	72/M		Grade II		
7	38/M	Pontomedullary angle	Grade I		Dysfunction VII - XII R
8	33/M	Os petrosum	Grade II	3×2×2 cm	12 y Strabism; later unsteady gait and decreased hearing AD. Sensory disturbance V branch 2 R; paresis m trapezius and sternocleidom. R; tongue deviating to R; dysdiadochokinesis and ataxia R
9	47/F	Behind os petrosum; extension to PS	Grade I	4×4×4 cm	
10	64/F	Cranial clivus; SS up to Monro	Grade I	4×4×6 cm	Hemiparesis R; disorder of speech; memory disturbance
11	72/F	Clivus; PS; Sphenoid and SCav	Grade I		3½ y Decreased visual acuity OD; 3 months amaurosis OD and decreased visual acuity OS. Quadrant-anopsia and L VI dysfunction
12	45/M	FM and FP			
13	60/M	Petroclival			
14	44/M	Foramen jugulare and in tympanic cavity; Clivus and SCav	Grade II		2 Months pain R neck/ear; dizziness; nausea; decreased hearing and tinnitus AD; diplopia. Sensory disturbance R mouth, VII dysfunction R and hearing AD<AS
15	16/M	FM; Petrosal bone	Grade II		2 y Diplopia when looking R; paraesthesias maxillar and mandibular R, diplopia. No cornea reflex R; hypaesthesia V branch 1-2-3 R

PS = parasellar; FM = fossa media; FP = fossa posterior; SS = suprasellar; SCav = sinus cavernosus; SSph = sinus sphenoidalis.

Intracranial tumour growth can be present. The origin has been reported to be in different bone parts (table 2). In eight cases an origin in the dura was described.

A pathological diagnosis was reported in 106 cases (table 2). Four subtypes of skull base chondrosarcomas were described in the literature: grade I, II, mesenchymal, and myxoid type. No further pathological grading was given for mesenchymal and myxoid subtypes. The figure lists the pathological subtypes and their age of occurrence.

The extension in dural tissue of the chondrosarcoma, confirmed through postmortem data (10 cases) or radiological imaging only, was explicitly mentioned in 30 cases. Of these, 10 were diagnosed as chondrosarcoma grade I,

three were diagnosed as grade II, 16 were diagnosed as mesenchymal type, and one as myxoid.

The usual methods of treatment were neurosurgery and conventional or proton radiotherapy.^{38 40 41} Stereotactic radiosurgery has been reported in only two cases.²² Adjuvant chemotherapy was scarcely mentioned.⁴²

A local recurrence rate of 53% after neurosurgical treatment (34 out of 64 cases) has been reported with clinical and radiological signs of regrowth after a mean interval of 32 months. In 40 cases information on the extent of the operation was reported: in 80% (32 cases) resection was subtotal and in 20% (eight cases) total.⁴⁻³⁷

Thirteen of the 15 Dutch patients were operated on. Two received proton radiotherapy and one conventional radiotherapy postoperatively. Recurrence occurred in 54% (seven cases). The mean time to recurrence was three years. Recurrence free survival rates at 2, 3, and 5 years were calculated at respectively 67%, 56%, and 43%.

Discussion

This review of the literature and of our own cases (in total 192) illustrates the very low frequency of occurrence of intracranial chondrosarcoma. The incompleteness of data in the case reports is of such an extent that the clinical picture had to be made using a substantially smaller part of the published cases.

For patient characteristics, no sex dominance existed. The mean age was 37 years (our own series 43 years). Chondrosarcoma occurred in both very young and old age groups, from 3 months to 76 years of age. The mesenchymal subtype showed a tendency to occur at a younger age (figure).

The signs and symptoms at the first manifestation of the tumour were mainly caused by oculomotor dysfunction, related to

Table 2 Clinical presentation, tumour location, and pathological subtypes of chondrosarcoma of the cranial base

		Cases	(%)
Symptoms/signs	Dysfunction of eye movement	37	51
	Headache	22	31
	Hearing loss, dizziness, and tinnitus	15	21
	Sensory disturbances of the face	15	21
	Decreased visual acuity	10	14
	Dysfonia	9	13
	Ataxia	8	11
	(Hemi)paresis	7	10
	Dysarthria/dysphagia	6	8
	Tongue paresis	6	8
	Facial palsy	6	8
	Nasal obstruction	3	4
	Decreased memory	3	4
	Nausea/vomiting	2	3
	Palatal mass	2	3
Bone parts	Os frontale	2	
	Os ethmoidale	3	
	Os sphenoidale	25	
	Os petrosum	47	
	Os parietale	13	
	Os occipitale or clivus	30	
	Origin in dura	8	
Subtypes	Grade I	54	51
	Grade II	12	11
	Grade III	0	
	Mesenchymal	32	30
	Myxoid	8	8

Table 1 continued

Invasion	Postoperative course
Dural and cerebral	Epilepsy and persisting VI dysfunction R; no recurrence after 10 y
Dural	Pain maxillar and mandibular L, decreased hearing AS; recurrence after 2 y (operated) Mild III/VI dysfunction; twice reoperated on; recurrence after 3 y: proton radiotherapy Persistent III dysfunction L 1 y After operation proton radiotherapy; no recurrence 3 y after operation
Dural	Dysarthria, L hemiparesis and abscess cerebral, epilepsy; 4 months later recurrence Recurrence after 8 y Recurrence after 2 y; death 19 y after diagnosis
Dural and cerebral	Three times reoperated on; postoperative V, VII, IX, and XI dysfunction; epilepsy Increased paresis and dysphasia No surgical treatment
Dural	Reoperation 1 y later, death 11 y after diagnosis Total hearing loss AD and VII dysfunction R; later stereotactical radiotherapy Decreased visual acuity OD and hearing AD; hypaesthesia R face; no recurrence 3 y after operation

the preferable location of chondrosarcoma in the petrosal part of the skull base. A more lateral localisation explains initial VIIIth nerve dysfunction. Cerebral lesions caused by intradural and intracerebral expansion were reported in 30 cases. Most cases were described as originating from the skull base (table 2), possibly from remnants of embryonal cartilage. However, a few were reported to originate from

dural tissue, suggestive of metaplasia of meningeal fibroblasts.²

On radiological examination, plain skull radiography showed only bone destruction and calcifications. Before 1980, angiography was used to evaluate the intracranial extension. Skull CT with intravenous contrast also disclosed bone destruction and calcification with more detailed demarcation of the tumour extension.

Skull MRI with administration of gadolinium DTPA resulted in even better tumour demarcation and visualisation of dural extension. The anatomical relations of the tumour with main vessels such as the carotid arteries and optic nerves are important guides for the extension of neurosurgical extirpation.³⁹

Bone biopsy was the main diagnostic tool. The pathological classification of the subtypes grade I - III is based on differences in characteristics such as nuclear size, cellularity, mitotic rate, and frequency of lacunae with multiple nuclei.⁴³ Our study of 177 patients in the literature and 15 Dutch patients disclosed no grade III chondrosarcoma. In the mesenchymal subtype, primitive spindle cells are present.⁴⁴ The myxoid type is composed of strings of rounded cells in a more or less myxoid matrix.⁴⁵


With the help of tumour markers such as vimentin, cytokeratin, and S100, the chondrosarcoma can be differentiated from chordoma. Chordomas lack vimentin immunoreactivity and chondrosarcomas fail to express cytokeratin. S-100 protein expression is present in both.⁴⁵

In the literature, mesenchymal and myxoid subtypes are classified separately from the conventional type I - III gradation. Lichtenstein and Bernstein considered the mesenchymal chondrosarcoma as a separate subtype.⁴⁴ Evans *et al* described a subgroup of grade III chondrosarcoma containing large areas exhibiting a spindle cell pattern, but with a mitotic rate well in excess of minimum criteria for grade III.⁴³ Although not explicitly mentioned, their description fits the definition of mesenchymal chondrosarcoma. It is our opinion that on the basis of this finding, at least the mesenchymal chondrosarcoma should be considered as a separate subtype.

Mesenchymal chondrosarcomas mainly occurred in the younger age group (10-30 years). The grade I chondrosarcoma had no clear age preference.

Very little data were available on the presence of dural invasion in skull base chondrosarcoma. The more malignant mesenchymal chondrosarcoma had a tendency towards a malign growth pattern which is illustrated by the relatively high frequency of extension into dural and cerebral tissue. In our own series (15 patients), dural invasion was present in five cases: three of these were grade I and two grade II chondrosarcoma.

Metastases of chondrosarcoma with a primary localisation in other parts of the body were found in 10% of grade II and 71% of grade III tumours.⁴³ Metastases from the skull base were mentioned by Hassounah *et al* in five



Age distribution for the different pathological subtypes of skull base chondrosarcomas.

Table 3 Results of radiotherapy in 108 patients with low grade chondrosarcoma of the cranial base, treated with charged particles

Reference	Patients	Follow up (months)	Particles	Dose (GyE)	LC-5 (%)	SV-5 (%)	Complication rate (%)
Castro <i>et al</i> ³⁸	27	4–191 (median 51)	Helium, neon	60–80 (median 66)	78	83	41 (77–86) 20 (87–92)
Munzenrider <i>et al</i> ¹⁶	81	2–187 (median 37)	Proton	62.8–77.4 (median 68.5)	91	94	13 (auditory) 12 (endocrine) 10 (brain) 6 (visual)

LC-5 = local control rate at 5 y; SV-5 = survival rate at 5 y; GyE = Gray equivalent.

of 50 patients.⁴ We were not able to find other cases with metastases.

Neurosurgery is one of the treatment options. In 53% of neurosurgically treated patients recurrence of the tumour was found (mean interval 32 months). The high recurrence rate is caused by partial resection due to the proximity of critical neuronal and vascular structures.

Several studies describe neurosurgical procedures and their results in treatment of intracranial chondrosarcoma.^{18 34 41} Up to nine different surgical approaches are possible, often combined or in stages. Macroscopical total resection was accomplished in 56%–67% of cases. Postoperative radiotherapy was given in 20%–44% of the patients.

Gay and Sekhar reported on 60 patients with low grade cranial base chondrosarcoma¹⁴ and chordoma.^{41 46} Fifty percent of these patients were treated previously elsewhere. All were operated on and total or near total resection was achieved in 67% of cases. Twenty per cent of the patients received radiotherapy postoperatively. They found a recurrence free survival rate at five years of 65%. The most frequent complication of the operation was leakage of CSF (30%). During follow up two patients (3%) died because of complications of radiotherapy and three (5%) because of systemic complications of surgery.

Risk of recurrence of tumour growth was found to be greater in patients who were already operated on elsewhere and in the case of only partial resection.

In general, an optimal tumour removal in one operation was advocated, because repeated surgical intervention has risks of tumour progression, development of scar tissue, and secondary spread of tumour cells.^{18 34 41}

Radiotherapy is another treatment for these tumours in the direct vicinity of essential structures. The anatomical close relation between the tumour and these critical normal structures limits the dose that can be delivered with conventional radiation treatment. Charged particle radiotherapy combined with three dimensional treatment planning results in superior dose distribution that allows delivery of high tumour dose with acceptable dose to the normal tissues.

Table 3 shows the results of the treatment of low grade chondrosarcoma of the skull base with combined proton and photon radiotherapy and with helium or neon radiotherapy.^{38 46} Local control rates of 78%–91% and survival rates of 83%–94%, both at five years, were achieved.

The treatment related morbidity ranged from 6% (visual complications) to 13% (auditory complications) in series of chondrosarcomas (81 cases) and chordomas (113 cases) of both the skull base and cervical spine.⁴⁶ Castro *et al* reported a decline in occurrence of all radiotherapy complications from 41% in the period before 1986 to 20% after 1986, demonstrating the impact of improved imaging and treatment planning techniques.³⁸ However, in the period 1977–92, five of 85 disease free patients (5%) died due to complications of therapy.

Despite the high rates of local success of charged particle radiotherapy and the fact that local recurrence always led to deterioration of the neurological condition, morbidity and mortality are considerable and should be taken into account when offering this treatment to patients.

Up to now, charged particle radiotherapy can only be carried out in highly specialised centres. It is our opinion however, that the role of radiotherapy in the treatment of skull base chondrosarcoma, possibly in combination with surgery, in the years to come will grow.

Skull base chondrosarcomas are tumours that should be considered in cases of unexplainable cranial nerve dysfunction and other associated symptoms. The reported interval between the first symptoms and moment of diagnosis was often long. In most reports important data in defining the natural history of a chondrosarcoma were missing. Also, pathological classification should be given in more detail. In comparison with neurosurgical, mostly partial, tumour resection, charged particle radiotherapy seems to give better follow up results in terms of five year survival. Both neurosurgery and radiotherapy can cause treatment related side effects that should influence therapy choice in the individual patient.

We are indebted to the "Dutch Committee on Bone Tumours" for providing the necessary data on the patients in the Netherlands and to G Freling for his comments on pathological classification.

- Cianfriglia F, Pompili A, Occhipinti E. Intracranial malignant cartilaginous tumours. Report of two cases and review of the literature. *Acta Neurochir Wien* 1978;45:163–75.
- Bahr AL, Gayler BW. Cranial chondrosarcomas. Report of four cases and review of the literature. *Radiology* 1977;124:151–6.
- Volpe NJ, Liebsch NJ, Munzenrider JE, *et al*. Neuro-ophthalmologic findings in chordoma and chondrosarcoma of the skull base. *Am J Ophthalmol* 1993;115:97–104.
- Hassounah M, Al-Mefty O, Akhtar M, *et al*. Primary cranial and intracranial chondrosarcoma: a survey. *Acta Neurochir Wien* 1985;78:123–32.
- Adegbite ABO, McQueen JD, Paine KWE, *et al*. Primary intracranial chondrosarcoma: a report of two cases. *Neurosurgery* 1985;17:490–4.

- 6 Cybulski GR, Russell EJ, D'Angelo CM, et al. Falcine chondrosarcoma: case report and literature review. *Neurosurgery* 1985;16:412-5.
- 7 Hertzanu Y, Mendelsohn DB, Davidge-Pitts K, et al. Chondrosarcoma of the head and neck-the value of computed tomography. *J Surg Oncol* 1985;28:97-102.
- 8 Kveton JE, Brackmann DE, Glasscock ME, et al. Chondrosarcoma of the skull base. *Otolaryngol Head Neck Surg* 1986;94:23-32.
- 9 Arruda WO, Cardoso MD, Pedrosa AA, et al. Intracranial chondrosarcoma: report of a case. *Arg Neuropsiquiatr* 1986;44:89-93.
- 10 Masuzawa T, Nakahara N, Saito K, et al. Parasellar chondrosarcoma. Case report. *Neurol Med Chir (Tokyo)* 1986;26:44-8.
- 11 Coltrera MD, Googe PB, Harist TJ, et al. Chondrosarcoma of the temporal bone. Diagnosis and treatment of 13 cases and review of the literature. *Cancer* 1986;58:2689-96.
- 12 Nokes SR, Dauito R, Murtagh FR, et al. Cranial mesenchymal chondrosarcoma. *Am J Neuroradiol* 1987;8:1137-8.
- 13 Lee Y, Van Tassel P, Raymond AK. Intracranial dural chondrosarcoma. *Am J Neuroradiol* 1988;9:1189-93.
- 14 Boorstein JM, Spizarny DL. Case report 476. *Skeletal Radiol* 1988;17:208-11.
- 15 Charachon R, Crouzet G, Vasdev A, et al. Tumours of the petrous apex: a clinical series. *J Neuroradiol* 1988;15:186-201.
- 16 Sindou M, Daher A, Vighetto A, et al. Parasellar chondrosarcoma: report of a case operated on through a pteriono-temporal approach and review of the literature. *Neurochirurgie* 1989;35:186-90.
- 17 Shuangshoti S, Kasantikul V. Primary intracranial mesenchymal chondrosarcoma. *J Laryngol Otol* 1989;103:545-9.
- 18 Nath-Sen C, Sekhar LN, Schramm VL, et al. Chordoma and chondrosarcoma of the cranial base: an 8-year experience. *Neurosurgery* 1989;25:931-41.
- 19 Oguro K, Nakahara N, Yamaguchi Y, et al. Chondrosarcoma of the posterior fossa. Case report. *Neurol Med Chir (Tokyo)* 1989;29:1030-8.
- 20 Greenblatt SH. Left occipital lobectomy and the preangular anatomy of reading. *Brain* 1990;38:576-95.
- 21 Miyamori T, Mizukoshi H, Yamano K, et al. Intracranial chondrosarcoma. Case report. *Neurol Med Chir (Tokyo)* 1990;30:263-7.
- 22 Kondziolka D, Lunsford LD, Flickinger JC. The role of radiosurgery in the management of chordoma and chondrosarcoma of the cranial base. *Neurosurgery* 1991;29:38-46.
- 23 Kretschmar HA, Eggert HR. Mesenchymal chondrosarcoma of the craniocervical junction. *Clin Neurol Neurosurg* 1990;92:343-7.
- 24 Koga H, Mukawa J, Miyagi K, et al. A case of chondrosarcoma in the cavernous sinus. *No Shinkei Geka* 1991;19:1167-70.
- 25 Salzman M, Scholtz H, Krist D, et al. Extraskelletal myxoid chondrosarcoma of the falx. *Neurosurgery* 1992;31:344-8.
- 26 Wojno KJ, Hruban RH, Garin-Chesa P, et al. Chondroid chordomas and low-grade chondrosarcomas of the craniospinal axis. An immunohistochemical analysis of 17 cases. *Am J Surg Pathol* 1992;16:1144-52.
- 27 Antonaci F, Sjaastad O. Hemispheric continua: a possible symptomatic case, due to mesenchymal tumor. *Funct Neurol* 1992;7:471-4.
- 28 Eavey RD, Janfaza P, Chapman PH, et al. Skull base dumbbell tumor: surgical experience with two adolescents. *Ann Otol Rhinol Laryngol* 1992;101:939-45.
- 29 Ruark DS, Schlehaider UK, Shah JP. Chondrosarcomas of the head and neck. *World J Surg* 1992;16:1010-6.
- 30 Çolak A, Berker M, Saglam S, et al. Chondrosarcoma of the temporal bone in an infant: case report and review of the literature. *Neurosurgery* 1992;31:956-7.
- 31 Bourgouin PM, Tampieri D, Robitaille Y, et al. Low-grade myxoid chondrosarcoma of the base of skull: CT, MR, and histopathology. *J Comput Assist Tomogr* 1992;16:268-73.
- 32 Makhmudov UB, Tcherekayev VA, Tanyashin SV. Transoral approach to tumors of the clivus: report of two cases. *Journal of Craniofacial Surgery* 1992;3:35-8.
- 33 Cho BK, Chi JG, Wang KC, et al. Intracranial mesenchymal chondrosarcoma: a case report and literature review. *Childs Nerv Syst* 1993;9:295-9.
- 34 Stapleton SR, Wilkins PR, Archer DJ, et al. Chondrosarcoma of the skull base: a series of eight cases. *Neurosurgery* 1993;32:348-56.
- 35 Stiglbauer R, Böhm G. Chondromatöse Tumoren in der MRT: Erscheinungsbild in Abhängigkeit von Lokalisation und Histopathologie. *Fortschritte auf dem Gebiete der Röntgenstrahlen und der Nuklear Medizin* 1993;158:147-53.
- 36 Volpe NJ, Lessell S. Remitting sixth nerve palsy in skull base tumors. *Arch Ophthalmol* 1993;111:1391-5.
- 37 Sato K, Kubota T, Yoshida K, et al. Intracranial extraskelletal myxoid chondrosarcoma with special reference to lamellar inclusions in the rough endoplasmic reticulum. *Acta Neuropathol* 1993;86:525-8.
- 38 Castro JR, Linstadt DE, Bahary JP, et al. Experience in charged particle irradiation of tumors of the skull base: 1977-92. *Int J Radiat Oncol Biol Phys* 1994;29:647-55.
- 39 Meyers SP, Hirsch WL, Curtin HD, et al. Chondrosarcoma of the skull base: MR imaging features. *Radiology* 1992;184:103-8.
- 40 Gay E, Sekhar LN, Rubinstein E, et al. Chordomas and chondrosarcomas of the cranial base: results and follow-up of 60 patients. *Neurosurgery* 1995;36:887-96.
- 41 Austin-Seymour M, Munzenrider J, Linggood R, et al. Fractionated proton radiation therapy of cranial and intracranial tumors. *Am J Clin Oncol* 1990;13:327-30.
- 42 Yoshimoto T, Sawamura Y, Ikeda J, et al. Successful chemoradiation therapy for high grade skull base chondrosarcoma in a child. *Childs Nerv Syst* 1995;11:250-3.
- 43 Evans HL, Ayala AG, Romsdahl MM. Prognostic factors in chondrosarcoma of bone. A clinicopathologic analysis with emphasis on histologic grading. *Cancer* 1977;40:818-31.
- 44 Lichtenstein L, Bernstein D. Unusual benign and malignant chondroid tumors of bone. A survey of some mesenchymal cartilage tumors and malignant chondroblastic tumors, including a few multicentric ones, as well as many atypical benign chondroblastomas and chondromyxoid fibromas. *Cancer* 1959;12:1142-57.
- 45 Brooks JJ, LiVolsi VA, Trojanowski JQ. Does chondroid chordoma exist? *Acta Neuropathol (Berl)* 1987;72:229-35.
- 46 Munzenrider JE, Liebsch NJ, Dubois W, et al. High dose fractionated combined proton and photon radiation therapy of chordomas and low-grade chondrosarcomas of the skull base and cervical spine. I. Results in adult patients. Abstract. *Radiation Oncology Biology Physics* 1991;21:16.